

Mushroom: Cultivation and Processing

Ashutosh kumar, Prof. Brajesh kumar and Mr. Karn Kumar

INTRODUCTION:

Mushrooms are one of the most loved food not only for its exotic taste but also for the benefits with which it comes. It can be consumed in various forms like fresh, pickled, dried, powdered, canned etc. Its farming has picked up a fast pace among contemporary entrepreneurs owing to its nutritional and medicinal benefits and low cost input with high output. Mushrooms are a fleshy fungi (Basidiomycota, Agaricomycetes) having a stem, cap and gills underneath the cap.

They can be edible, wild and some of them can be toxic too. It contains more than 90% water and less than 1% fat, loaded with Vitamin B, copper and selenium and low in sodium.

Usually vegetables, milk and other food products are fortified with Vitamin D by irradiation or direct addition but mushrooms are unique in this sense because they are naturally a rich source of Vitamin D which otherwise is procured from animals or poultry.

The reason being that it contains copious amount of plant sterol “Ergosterol”.

It is a precursor of Vitamin D which when stimulated by sunlight or artificial lightening source converts to Vitamin D.

Table 1- presents the details of nutritional value of *Agaricus bisporus* (White Button Mushrooms).

Table 1: Nutritional Value of *Agaricus bisporus* (White Button Mushrooms) per 100 gm

Nutritional Value per 100 g (3.5 oz.)	
Energy	113 kJ (27 kcal)
Water	92.45 g
Carbohydrates	4.1 g
Fat	0.1 g
Protein	2.5 g
Thiamine (Vit B1)	0.1 mg (9%)
Riboflavin (Vit B2)	0.5 mg (42%)
Niacin (Vit B2)	3.8 mg (25%)
Pantothenic Acid (B5)	1.5 mg (30%)
Vitamin C	0 mg (0%)
Calcium	18 mg (2%)
Phosphorous	120 mg (17%)
Potassium	448 mg (10%)
Sodium	6 mg (0%)
Zinc	1.1 mg (12%)
Vitamin D (D2 + D3)	0.2 µg
Sugar	1.98 g

**Percentages are relative to US
recommendations for adults.
Source: USDA Nutrient Database**

¹Ashutosh kumar, (Research scholar) S.D.J.P.G. College, Chandeswar, Azamgarh, (UP).

²Prof. Brajesh kumar, (HOD Plant Pathology) S.D.J.P.G. College, Chandeswar, Azamgarh, (UP).

³Mr. Karn Kumar, B.Sc (Ag), M.M.P.G.C., Kalakankar, Pratapgarh (U.P.).

TYPES OF MUSHROOMS

Mushrooms are easily cultivable in hilly regions due to abundant moisture but can also be grown in artificial environment with proper temperature and humidity control. Varieties must be identified thoroughly as some of them might cause food poisoning or allergy upon consumption. Some of the major varieties consumed in India are as follows:

Button Mushroom

Button mushroom (*Agaricus bisporus*) belongs to Class Basidiomycetes and Family Agaricaceae and is native to Europe and North America. It is of two types white and brown, out of which white button mushroom- is commonly grown in India.



According to ICAR - Directorate of Mushroom Research, this variety contributes more than 85% to mushroom production.

- It is the most relished variety used in eateries and households.

Shiitake Mushroom

Shiitake Mushrooms are native to East Asia and are highly consumed in Asian

countries. They readily grow on wood of deciduous and hard wood trees such as Oak, Chestnut, and Maple etc. and require moist and warm climate. In rare cases they may cause allergic reaction like itching but can be eliminated by thorough cooking. These are used in Asian cuisines and traditional medicines.

Oyster Mushroom

Oyster Mushrooms (*Pleurotus ostreatus*) belongs to *Pleurotus* species. It is known as “Dhingri” in India and has fan or oyster shaped cap. They grow easily on decaying wood or straw.



Paddy Straw Mushroom

Volvariella volvacea belongs to division Basidiomycota. It is usually grown on Rice straw bed and is used extensively in Asian Cuisines.

CULTIVATION

The basic requirements for mushroom cultivation are manure/compost, spawns, right temperature and humidity. Favorable growing conditions involve 80%- 90% of relative humidity, ample ventilation, a temperature range of 20-28⁰ C during spawn run and 12-

18⁰ C for reproductive growth. Initially for a week temperature must be maintained at 23 ± 2⁰ C and then it can be reduced to 16 ± 2⁰ C for subsequent weeks. The CO₂ concentration should be 0.08-0.15 % [13]. If the above stated conditions are maintained appropriately the pin heads start to appear within few days and progressively mature into button stage. Apart from these insecticides, nutritional supplements like nitrogen, vermiculite, water are also required for a healthy harvest. The following steps are to be followed for mushroom cultivation:

Compost Preparation

The compost (synthetic or natural) used for mushroom growth usually comprises of wheat straws, horse manure, poultry manure, rice bran, gypsum etc.

Utmost care is taken to protect the raw compost against rain (or external moisture, as it might introduce undesirable microbes. The chopped wheat straws or rice bran are mixed with horse dung, sprinkled with water and are heaped in a pile to allow fermentation. The fermentation process along with heat development breaks down the chemical compounds in small components. Frequent turnings and watering is done at a specific interval so as to avoid the drying up of compost. Gypsum is sometimes added to the compost to reduce greasiness and allow more aeration. Within

15 to 20 days the compost gets all set to be used as bed, it is then spread onto wooden trays and sowed with spawns.

Spawning

Spawns refers to the mycelium carefully propagated on agars or grains. Spawning is a process of sowing or mixing spawns in compost. Although mushroom produces spores which acts as a seed for further propagation but are not used generally due to uncertain germination and growth. The spawns are thoroughly mixed with the compost, are covered with newspaper and is watered sufficiently to maintain the moisture. Throughout the cultivation period humidity is kept high to avoid loss of moisture. Gradually they grow into white cottony mycelium growth.

Casing

Casing is a kind of sterilized soil or dressing containing cow manure which is spread onto the spawn mixed compost. It is applied when the mycelium growth commences on the compost surface. After 15 to 20 days of its application mushroom head or pins start becoming visible on the surface. They are allowed to mature for a specific time period and are harvested before opening of the cap. Mushrooms with opened cap (looks like an umbrella after opening of cap) are undesirable and are considered of menial quality.

HARVESTING

Harvesting is done by plucking them from soil using hands or the heads are chopped off using knife. The harvested mushrooms are then subjected to primary processing.

PROCESSING

Mushroom are very fragile and have a short shelf life, unless consumed fresh. At ambient temperature they lose their freshness within a day and deteriorates rapidly if not processed or refrigerated. They also tend to brown due to presence of compound Tyrosinase. It converts monophenols to diphenols which in turn are oxidized to quinones resulting in the formation of insoluble brown pigment called Melanin.

Initial processing involves washing mushrooms to remove adhering soil or compost and blanching them for few minutes to inactivate the enzymes. In order to prevent discoloration they are treated with brine, salt or citric acid prior to canning or packaging. Following are the common processing and preservation methods used:

Drying

Drying or Dehydration is the oldest and the basic processing method for various food products. Moisture is the most suitable medium for the microbial growth and

propagation, hence its removal will cause the microbial activity to cease or become gradual. Mushrooms can be dried either by sun drying or by mechanical drying. Sun drying is the cheapest and popular method but it produces a much darker product [19]. Mechanical drying is rapid and is of various types like Tray drying, Freeze drying, Vacuum drying, Microwave oven drying, Air drying etc. Dried mushrooms can be rehydrated and are used in soups, stews, pickles etc.

Freezing

As mushrooms contain more than 90% water hence freezing is the most suitable method for preservation. They are subjected to various pretreatments to minimize unfavorable effects upon freezing. According to Czapski and Szudyga (2000) colour of *Agaricus bisporus* is better when treated with metabisulphites along with blanching. Blast freezing method is commonly employed at temperature -25°C to -30°C (Sobkowska and Wozniak 1974, Czapski and Szudyga 2000). According to Kondratowicz and Kowalko (2000) Cryogenic freezing extends mushroom's shelf life up to one year when used at -80°C to -100°C for 5-6 minutes.

Sterilization

Sterilization of mushrooms can be done by using chemicals, steam or by irradiation. According to Kashif Akram and Joong-Ho Kwon (2010) the shelf life of

mushrooms can be extended by applying a radiation dose of 1 – 3 kGy.

Canning


Canning involves preservation in brine, vinegar, oil or marinades. Freshly harvested mushrooms are utilized for canning purpose. They are cleaned, graded, blanched and then filled into cans along with brine or vinegar followed by lidding. The cans are then exhausted to remove air, heat sterilized, cooled, labelled and packaged for storage or consumption.

Pickling

Pickling is an age old method which utilizes spice, salt, vinegar and oil as the basic ingredients for food preservation. Pickled mushrooms are made using spices such as turmeric, red chilli, garlic, clove along with salt and oil. Pickling induces fermentation which generates a mild flavor.

DISCUSSION

Mushrooms are adaptable fungi which yields lots of nutritional and health benefits. It can be grown on decomposed matter easily with good yield. They are natural source of vitamin D. There are lots of varieties including edible as well as wild. The commonly consumed varieties are Button mushroom (*Agaricus bisporus*), Shiitake mushroom, Oyster mushroom and Paddy Straw mushroom. Mushroom cultivation require 80-90% relative humidity and temperature range of 16° C - 23° C. The cultivation starts with compost preparation followed by spawning, casing, harvesting and processing. Mushroom being fragile and perishable commodity requires processing after harvesting. The processing can be accomplished in various ways such as drying, dehydration, freeze drying, pickling, freezing, canning, sterilization, direct packaging etc.



(A). Button Mushroom



(B). Oyster Mushroom



Paddy straw Mushroom

(C). Paddy straw Mushroom



Shitake Mushroom

(D). Shitake Mushroom

CONCLUSION

Mushrooms are one of the most popular and versatile gift of nature. It can be mixed into any food preparations or can be processed to give a new product. A lot of mushroom products are currently available in market such as mushroom pickle, seasonings, beverages, extracts, dried and canned mushrooms, mushroom supplements, cosmetics etc. Apart from the mushroom food products many innovative products are emerging in other industries as well such as mushroom based building materials, medicines, mycelium based platforms, biodegradable packaging, mycelium based leather etc. Mushrooms are easy to cultivate, have quick growth and nil carbon emission and waste generation. The fungi is a good source of income generation for the growers and also provides additional benefits through its processing.

Hence mushrooms hold a bright future

in every aspect owing to its diverse properties.

REFERENCES

1. Ainaworth , G. C., Sussman, A. S. and Sparrow, F. K. (1973).The fungi, vol. 4.B. Academic press, New Yourk.
2. Anonymous (2004). A.I.C.N.I.P. Work Shop, Raipur Cente, Department of Plant Pathology, IGAU, Raipur.
3. Antwi,E.(1995).Influence of saw dust supplemented with farm and industrial wastes on the growth, yield and mineral composition of *Pleurotus ostreatus* (Jacq. ex.fr.) Kummos.Munoz, Nueva Ecija. 150 leave.
4. Awasthi, S. K. (1992). Physico-chemical needs for high mycelia yield of *Pleurotus flabellatus* (Berk. and Br.) Sacc., an edible mushroom. National Academy Science letters,**15** (1): 1-2.
5. Badcock, E.C. (1943). Methods of

- obtaining fructification of wood rotting fungi in culture. *Trans. Br. Mycol. Soc.* **26**: 127-132.
6. Bahukhandi, D. (1986). Studies on some edible *Pleurotus species*, with particular reference to development yielding strains and improvement in production technology. Ph.D. thesis, Department of Botany, University of Garwal, Srinagar. 146p.
 7. Bahukhandi, D. and Munjal, R. L. (1989). Cultivation of *Pleurotus species* on different agricultural residues. *Indian PHYTOPATH.* **42**: 492-495.
 8. Bahukhandi, D. ;Bhal, N. Mahar, M. J. (1991). Colony characters of some strain of *Agaricus bitorquis* (Quel) Sacc. *Mushroom Science XIII, Vol. 1. Proceeding of the 13th International congress on the science and cultivation of edible fungi.* Bublin, Iris Republic, 1-6 Sept. pp.111-114.
 9. Bahukhandi, D. ;Bhal, N. Mahar, M. J. (1991). Colony characters of some strain of *Agaricus bitorquis* (Quel) Sacc. *Mushroom Science XIII, Vol. 1. Proceeding of the 13th International congress on the science and cultivation of edible fungi.* Bublin, Iris Republic, 1-6 Sept. pp.111-114.
 10. Chandra, S., Singh, A.K., Bhat, M.N. and Kumar, S. (1998). Yield performancr of *Pleurotus sajor caju* on some selected substrates in North-Eastern region of India. *Mush.Res.* **7**(2):79-80.
 11. Chandrasekhar, B.S. and Singh, S. (2006). Strain improvement in Oyster mushroom (*Pleurotus sajor caju*) through mutation breeding. *Mush.Res.*, **15**(1): 19-22.
 12. Dugger, B.M. (1905). Some principles in mushroom growing and mushroom spawn making. *Bull, U.S.Dept. Agric.*, **85**:1-60.
 13. Hazarika, D.K. (1998). Effect of substrates and supplements on yield of Oyster mushroom (*Pleurotus sajor caju*). *Mush. Res.*, **7**(1): 47-48.