

Beyond Orange: Unlocking the Genetic and Nutritional Diversity of Carrot

Veer Bhan singh, Manasi singh , Dr Muzeev Ahmad

Introduction:

Carrot (*Daucus carota* L.) is one of the most familiar root vegetables in the world, commonly recognized by its bright orange colour. However, the biological diversity of carrot extends far beyond orange roots. Carrot germplasm includes **orange, purple, black, red, yellow and white roots**, each representing different combinations of pigments, phytochemicals and genetic backgrounds. This diversity makes carrot an important crop not only for vegetable production but also for **nutritional improvement, functional foods and modern**

plant breeding. The nutritional characteristics of carrot are strongly influenced by genotype. Studies have demonstrated variation among carrot cultivars in carotenoids, phenolic compounds, anthocyanins, vitamins and antioxidant capacity. Therefore, the colour of a carrot can provide an initial visual indication of underlying biochemical diversity, although colour alone cannot fully describe its nutritional composition.

A Rainbow Beneath the Soil

The familiar orange carrot is rich in carotenoids, particularly α -carotene and β -

Table 1. Colour diversity and major nutritional characteristics of carrot

Root colour	Major associated compounds	Breeding/nutritional significance
Orange	α -carotene, β -carotene and other carotenoids	Important source of provitamin-A carotenoids
Purple/black	Anthocyanins, phenolic compounds	Potential source of antioxidant phytochemicals and natural colourants
Red	Carotenoids; lycopene in some genetic backgrounds	Provides additional pigment and phytochemical diversity
Yellow	Xanthophylls and other carotenoids	Useful source of diverse carotenoid profiles
White	Lower coloured-pigment concentration; other phytochemicals	Valuable genetic material for studying pigment pathways

Source: Synthesized from Arscott and Tanumihardjo (2010), Nicolle et al. (2004), and Iorizzo et al. (2020).

Veer Bhan singh, Manasi singh , Dr Muzeev Ahmad

Department of Genetics and plant breeding,

Acharya Narendra Deva University of Agriculture & Technology, Ayodhya, 224 229 (U.P.)

carotene. β -carotene is nutritionally important because it can be converted by the human body into vitamin A. Purple and black carrots, in contrast, are characterized by the accumulation of **anthocyanins**, while yellow carrots possess distinctive carotenoid profiles. Red carrots may contain different

combinations of carotenoids, including lycopene in particular genetic backgrounds. White carrots generally contain relatively low levels of coloured carotenoids. Consequently, carrot colour represents more than an aesthetic trait—it reflects underlying biochemical and genetic variation.

Colour diversity and major nutritional characteristics of carrot



Figure 1- Colour diversity and major nutritional characteristics of carrot

Beyond Orange: Carrot Diversity and Breeding Opportunities




Figure 1. Conceptual framework linking carrot colour and genetic diversity with nutritional traits and modern breeding strategies. The figure is an original synthesis based on published research

Genetic Diversity: The Raw Material of Breeding

Genetic diversity is the foundation of crop improvement. In carrot, variation occurs for numerous economically and nutritionally important traits, including root colour, shape, length, diameter, weight, carotenoid concentration, anthocyanin content, maturity, flavour, disease response and environmental adaptation. The domestication history of carrot has involved selection for root characteristics and other desirable traits. Transcriptome-based research comparing cultivated and wild carrots identified thousands of single-nucleotide polymorphisms and genes showing evidence of selection, demonstrating the genetic complexity underlying carrot domestication.

For breeders, this diversity provides opportunities to combine favourable alleles

from different genetic backgrounds rather than relying solely on existing commercial cultivars.

Orange Carrot: A Natural Source of Provitamin A

Orange carrots owe their characteristic colour primarily to the accumulation of carotenoids. Among these, β -carotene and α -carotene are particularly important because they possess provitamin-A activity. Historical breeding efforts have successfully increased provitamin-A carotenoid concentrations in carrot. This demonstrates that nutritional composition can be treated as a breeding objective alongside yield, shape and market quality. The availability of the carrot genome

has further improved understanding of the molecular mechanisms controlling carotenoid accumulation. The chromosome-scale carrot genome identified gene families associated with pigment accumulation and provided an important foundation for genomics-assisted breeding.

Purple and Black Carrots: The Anthocyanin Treasure

Purple and black carrots are particularly valuable because their colour is associated with **anthocyanin accumulation**. Anthocyanins are water-soluble flavonoid pigments that contribute to the red, purple and blue colours of many plants. Research has shown considerable variation among purple carrot germplasm for both **total anthocyanin concentration and pigment composition**. The relative proportions of acylated and non-acylated anthocyanins can also differ among

genetic stocks. Such differences are important because pigment structure can influence stability and potential application as a natural food colourant. Importantly, anthocyanin pigmentation is genetically complex. Molecular studies have identified numerous loci and QTLs associated with anthocyanin accumulation in carrot. This information can assist breeders in developing cultivars with predictable pigment characteristics.

Nutritional Quality Is Controlled by Both Genes and Environment

Carrot nutritional quality is not determined by genetics alone. **Genotype × environment interactions, mineral nutrition, maturity and production conditions** can influence the concentration of bioactive compounds. For example, research evaluating different carrot cultivars demonstrated that both genetic background and mineral supply

Table 2. Major breeding targets for next-generation carrot cultivars

Target	Important traits	Potential breeding strategy
Nutritional enhancement	β -carotene, α -carotene, anthocyanins	Germplasm screening and marker-assisted selection
Root productivity	Root weight, diameter, length and marketable yield	Conventional selection and genomic selection
Colour diversity	Orange, purple, red, yellow and white	Phenotypic selection and pigment profiling
Functional-food potential	Anthocyanins, carotenoids and phenolics	Biochemical screening
Climate resilience	Heat, drought and disease tolerance	Stress screening + molecular breeding
Consumer acceptance	Flavour, texture, shape and uniformity	Multi-trait selection

influenced carotenoids, vitamin C, phenolic acids and antioxidant capacity. This indicates that nutritional improvement should integrate **genotype selection with appropriate crop management.**

The Genome Is Changing Carrot Breeding

The publication of a high-quality carrot genome assembly marked an important milestone in carrot genetics. Genome information has provided researchers with opportunities to investigate the genetic basis of **pigmentation, root development, flavour, flowering and stress responses.**

Modern carrot improvement can now combine conventional breeding with:

- ☞ Molecular markers
- ☞ Quantitative trait locus (QTL) analysis
- ☞ Genome-wide association studies
- ☞ Transcriptomics
- ☞ High-throughput phenotyping
- ☞ Genomic selection

☞ Candidate-gene analysis

☞ Genome editing

These technologies can shorten the path from identifying useful genetic variation to incorporating it into improved breeding populations. Reviews of carrot research indicate that genomics and high-throughput sequencing are increasingly contributing to nutritional studies, resistance breeding and cultivar development.

From Germplasm to Functional Carrot

The future of carrot breeding is likely to involve **multi-trait improvement** rather than selection for yield alone. A successful cultivar should ideally combine high productivity with nutritional quality, attractive appearance, desirable flavour, storage ability and adaptation to local environments. For example, a breeding programme could aim to develop an orange carrot with enhanced provitamin-A carotenoids while maintaining

Table 3. Conventional versus modern approaches in carrot improvement

Feature	Conventional breeding	Modern breeding
Selection basis	Visible phenotype	Phenotype + molecular information
Nutritional traits	Laboratory screening	High-throughput biochemical and genomic screening
Genetic information	Limited	Genome-wide information
QTL identification	Difficult and time-consuming	Marker-based identification
Selection efficiency	Depends strongly on phenotype	Can use markers linked with target traits
Potential for precision	Moderate	Higher
Major advantage	Simple and established	Efficient exploitation of complex genetic variation

good root shape and yield. Another programme could target purple carrots with high concentrations of stable anthocyanins for fresh consumption or natural food-colour applications.

Such approaches transform carrot from a conventional vegetable into a potential **functional-food crop**.

Conservation of Carrot Diversity: An Essential Investment

Genetic improvement cannot progress without genetic resources. Wild relatives, traditional varieties, landraces and older cultivars may contain alleles that are absent or rare in modern commercial populations. Conservation of carrot germplasm therefore serves two purposes. First, it protects biodiversity for future generations. Second, it provides breeders with a reservoir of genetic variation for addressing emerging challenges such as **climate stress, new diseases, changing consumer preferences and nutritional security**.

The study of carrot domestication has already demonstrated the value of comparing cultivated and wild genetic resources to understand the evolutionary changes that shaped the crop.

What Lies Beyond Orange?

The future of carrot is not simply about producing more roots—it is about producing **better and more diverse roots**. The

combination of conventional breeding, nutritional profiling and molecular genetics offers an opportunity to develop cultivars tailored for specific nutritional and agricultural purposes.

Imagine carrots bred specifically for:

- ⇒ **High provitamin A** → improved nutritional value
- ⇒ **High anthocyanin** → functional-food and natural-colour applications
- ⇒ **High carotenoid diversity** → broader phytochemical profiles
- ⇒ **Climate resilience** → stable production under environmental stress
- ⇒ **Superior root quality** → better consumer acceptance and market value

This approach places nutrition and resilience alongside yield as central objectives of crop improvement.

Conclusion

Carrot is a striking example of how genetic diversity can translate into nutritional diversity. The familiar orange root represents only one part of the crop's biological spectrum. Purple, black, red, yellow and white carrots contain different combinations of pigments and phytochemicals, providing valuable resources for both nutrition and breeding. The integration of germplasm conservation, phenotypic evaluation, biochemical screening, genomics and molecular breeding can accelerate the development of carrots with

improved nutritional value, productivity and resilience. The carrot of the future may therefore be defined not by a single colour, but by a portfolio of traits designed to meet the nutritional and agricultural needs of a changing world.

Beyond orange lies an entire spectrum of genetic possibilities—and that spectrum may hold the key to the next generation of nutritionally enhanced carrots.

References

1. Arscott, S. A., & Tanumihardjo, S. A. (2010). Carrots of many colors provide basic nutrition and bioavailable phytochemicals acting as a functional food. *Comprehensive Reviews in Food Science and Food Safety*, 9(2), 223–239. <https://doi.org/10.1111/j.1541-4337.2009.00103.x>
2. Cavagnaro, P. F., Bannoud, F., Iorizzo, M., Senalik, D., Ellison, S. L., & Simon, P. W. (2019). Carrot anthocyanins: Nutrition, diversity and genetics. *Acta Horticulturae*, 1264, 101–106. <https://doi.org/10.17660/ActaHortic.2019.1264.11>
3. Iorizzo, M., Ellison, S. L., Senalik, D., Zeng, P., Satapoomin, P., Huang, J., Bowman, M., Iovene, M., Sanseverino, W., Cavagnaro, P. F., Yildiz, M., Macko-Podgórn, A., Moranska, E., Grzebelus, E., Grzebelus, D., Ashrafi, H., Zheng, Z., Cheng, S., Spooner, D. M., Van Deynze, A., & Simon, P. W. (2016). A high-quality carrot genome assembly provides new insights into carotenoid accumulation and asterid genome evolution. *Nature Genetics*, 48(6), 657–666. <https://doi.org/10.1038/ng.3565>
4. Iorizzo, M., Curaba, J., Pottorff, M., Ferruzzi, M. G., Simon, P. W., & Cavagnaro, P. F. (2020). Carrot anthocyanins genetics and genomics: Status and perspectives to improve its application for the food colorant industry. *Genes*, 11(8), 906. <https://doi.org/10.3390/genes11080906>
5. Nicolle, C., Simon, G., Rock, E., Amouroux, P., & Rémésy, C. (2004). Genetic variability influences carotenoid, vitamin, phenolic, and mineral content in white, yellow, purple, orange, and dark-orange carrot cultivars. *Journal of the American Society for Horticultural Science*, 129(4), 523–529. <https://doi.org/10.21273/JASHS.129.4.0523>
6. Rong, J., Lammers, Y., Strasburg, J. L., Schidlo, N. S., Ariyurek, Y., de Jong, T. J., Klinkhamer, P. G. L., Smulders, M. J. M., & Vrieling, K. (2014). New

insights into domestication of carrot from root transcriptome analyses. *BMC Genomics*, 15, 895.
<https://doi.org/10.1186/1471-2164-15-895>

7. Singh, D. P., Beloy, J., McInerney, J. K., & Day, L. (2012). Impact of boron, calcium and genetic factors on vitamin C, carotenoids, phenolic acids, anthocyanins and antioxidant capacity of carrots (*Daucus carota*). *Food Chemistry*, 132(3), 1161–1170.
<https://doi.org/10.1016/j.foodchem.2011.11.045>

8. Zhang, Y., et al. (2019). Advances in research on the carrot, an important root vegetable in the Apiaceae family. *Horticulture Research*, 6, 69.
<https://doi.org/10.1038/s41438-019-0150-6>

