

Integrated Nutrient and Water Management for Sustainable AgricultureKiran N. More¹, Apeksha M. Suryawanshi², Shrikrishna B. Narale³**Introduction:**

Agriculture depends heavily on the availability of essential plant nutrients and adequate water. However, inefficient use of fertilizers and irrigation water can increase production costs, reduce soil fertility and contribute to environmental degradation. Declining groundwater levels, nutrient imbalance, soil degradation, irregular rainfall and increasing climate variability have further increased the need for efficient resource management. **Integrated Nutrient and Water**

Management (INWM) provides a practical approach to maintaining crop productivity while conserving soil, water and other natural resources. It involves the efficient and balanced management of organic and inorganic nutrient sources along with scientific water management practices to ensure that crops receive nutrients and water in the right quantity, at the right time and in the right place.

Concept of Integrated Nutrient and Water Management

Integrated Nutrient and Water Management is based on the principle that nutrient and water management should not be considered as separate activities. Availability of water influences nutrient uptake, while nutrient availability affects crop growth and water-use efficiency. Adequate soil moisture is necessary for nutrient movement and absorption by plant roots, whereas balanced nutrition helps crops develop stronger root systems and utilize available water more efficiently.

Integrated nutrient management combines chemical fertilizers with organic manures, compost, crop residues, green manures, biofertilizers and other locally available nutrient sources. Water management involves efficient irrigation, rainwater harvesting, moisture conservation, drainage management and scheduling irrigation according to crop demand. The integration of these practices can improve nutrient-use efficiency, water-use efficiency, crop productivity and soil health.

Kiran N. More¹, Apeksha M. Suryawanshi², Shrikrishna B. Narale³

¹ Ph.D. Scholar, Department of Agronomy, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth (DBSKKV), Dapoli, Maharashtra, India.

² Department of Animal Husbandry and Dairy Science (AHDS), Mahatma Phule Krishi Vidyapeeth (MPKV), Rahuri, Maharashtra, India.

³ Ph.D. Scholar, Department of Biochemistry, Mahatma Phule Krishi Vidyapeeth (MPKV), Rahuri, Maharashtra, India.

Importance of Nutrient Management

Plant nutrients are essential for growth, development and yield formation. Nitrogen, phosphorus and potassium are required in relatively large quantities, while secondary and micronutrients are also essential for balanced crop growth. Continuous cultivation and excessive use of particular fertilizers can create nutrient imbalances and gradually reduce soil fertility.

Integrated nutrient management aims to maintain an adequate and balanced supply of nutrients while minimizing nutrient losses. Soil testing is an important first step because it provides information about existing nutrient availability and helps determine fertilizer requirements. Fertilizer recommendations should consider soil fertility, crop requirement, expected yield and previous crop history.

Organic sources such as farmyard manure, compost and green manure can improve soil structure, microbial activity and organic matter. Chemical fertilizers provide nutrients in readily available forms and can meet crop requirements rapidly. Combining organic and inorganic sources can therefore provide both immediate nutrient availability and long-term improvement in soil fertility.

Organic Sources and Biofertilizers

Organic manures play an important role in sustainable nutrient management. Farmyard manure, compost, poultry manure,

crop residues and green manure can add organic matter and nutrients to soil. Regular addition of organic materials improves soil aggregation, water-holding capacity and biological activity.

Green manuring with suitable legume crops can contribute organic matter and biologically fixed nitrogen to the soil. Crop residues can also be retained or incorporated instead of being burned, helping recycle nutrients and improve soil health.

Biofertilizers containing beneficial microorganisms can contribute to nutrient availability through biological nitrogen fixation, phosphate solubilization and other processes. Their effectiveness depends on soil conditions, crop species, environmental conditions and appropriate application methods. Therefore, biofertilizers should be considered as a component of an integrated nutrient-management programme rather than a complete replacement for all fertilizers.

Balanced Fertilization

Balanced fertilization means supplying all essential nutrients in appropriate proportions according to crop requirements. Excessive application of nitrogen, particularly without adequate phosphorus, potassium and micronutrients, can result in poor nutrient-use efficiency and environmental losses.

The **4R nutrient stewardship principle—right source, right rate, right time

and right place—**provides a useful framework for improving fertilizer efficiency. Nitrogen fertilizers can be applied in split doses according to crop demand, while phosphorus and potassium can be managed according to soil-test recommendations. Placement of fertilizers near the active root zone can also improve nutrient availability and reduce losses.

Site-specific nutrient management and precision fertilizer application can further improve nutrient-use efficiency by considering spatial variability within fields.

Efficient Water Management

Water is one of the most critical resources for crop production, but its availability is becoming increasingly uncertain in many agricultural regions. Excessive irrigation can cause waterlogging, salinity, nutrient leaching and groundwater depletion, whereas inadequate irrigation can cause crop stress and yield reduction.

Efficient water management requires irrigation to be scheduled according to crop growth stage, soil moisture, weather conditions and water availability. Irrigation should be provided when the crop requires water rather than following a fixed schedule.

Methods such as drip and sprinkler irrigation can improve water-use efficiency under suitable conditions. Drip irrigation delivers water close to the plant root zone,

reducing evaporation and runoff. Sprinkler irrigation can provide relatively uniform water distribution and is suitable for many field and horticultural crops.

Soil-Moisture Conservation

Improving the soil's ability to retain water is an important component of sustainable water management. Practices such as mulching, residue retention, conservation tillage, contour farming and incorporation of organic matter can reduce evaporation and improve soil-water storage.

Crop residues left on the soil surface can protect the soil from direct solar radiation and reduce moisture loss. Organic matter improves soil structure and increases the capacity of soil to hold water. Conservation tillage also reduces soil disturbance and can help conserve moisture in suitable production systems.

Rainwater harvesting through farm ponds, field bunds and other water-conservation structures can provide additional water during periods of rainfall deficiency. In rainfed agriculture, capturing and conserving rainfall can be particularly important for stabilizing crop production.

Integration of Nutrient and Water Management

Nutrient and water management become more effective when they are planned together. Adequate moisture is required for

nutrient dissolution, movement and uptake, while balanced nutrition supports root development and enables crops to utilize water more effectively.

For example, excessive irrigation after fertilizer application can increase nutrient leaching, particularly for mobile nutrients such as nitrate. On the other hand, inadequate soil moisture can restrict nutrient uptake even when nutrients are present in the soil. Therefore, irrigation and fertilizer application should be synchronized with crop growth stages and environmental conditions.

and irrigation should be avoided when soil moisture is already adequate.

Water-Smart Fertilizer Application

The timing of fertilizer application is closely associated with water availability. Fertilizers should be applied when adequate soil moisture is available for nutrient dissolution and root uptake. However, excessive irrigation immediately after fertilizer application can increase nutrient losses.

Split application of nitrogen is particularly useful in crops with prolonged nutrient requirements. Applying nitrogen in

Table 1. Major Practices in Integrated Nutrient and Water Management

Practice	Major benefit
Soil testing	Provides a scientific basis for fertilizer recommendations
Balanced fertilization	Improves nutrient-use efficiency and crop growth
Farmyard manure and compost	Improves soil organic matter and water-holding capacity
Green manuring	Adds organic matter and biologically fixed nitrogen
Crop-residue management	Recycles nutrients and conserves soil moisture
Biofertilizers	Supports biological nutrient availability
Split fertilizer application	Synchronizes nutrient supply with crop demand
Drip irrigation	Improves water-use efficiency
Sprinkler irrigation	Provides efficient and uniform water application
Mulching	Reduces evaporation and conserves soil moisture
Rainwater harvesting	Stores water for use during dry periods
Conservation tillage	Reduces soil disturbance and improves moisture conservation
Soil-moisture monitoring	Helps determine appropriate irrigation timing
Integrated nutrient-water scheduling	Reduces losses and improves crop productivity

Integrated management can also reduce the risk of nutrient losses during heavy rainfall. Fertilizers should preferably be applied when extreme rainfall is not expected,

smaller doses during active crop-growth stages can improve synchronization between nutrient supply and crop demand. In irrigated agriculture, fertigation can further integrate

water and nutrient application by supplying soluble fertilizers through irrigation systems.

Fertigation is particularly useful with drip irrigation because nutrients can be delivered directly to the root zone. When properly managed, it can improve nutrient-use efficiency and reduce losses. However, fertilizer concentration, irrigation water quality and system maintenance must be carefully managed.

Role of Climate-Smart Practices

Climate variability has made integrated nutrient and water management increasingly important. Drought, heat waves, intense rainfall and changing rainfall patterns can affect nutrient availability and water requirements. Climate-smart management practices such as drought-tolerant varieties, conservation agriculture, mulching, efficient irrigation, rainwater harvesting and improved nutrient management can reduce production risks.

Improving soil organic matter is particularly important because soils rich in organic matter generally have better structure and water-holding capacity. Such soils can provide crops with greater resilience during short periods of water stress. Similarly, balanced nutrition can improve plant vigour and help crops recover from environmental stress.

Role of Precision Agriculture

Modern technologies can make integrated nutrient and water management more precise. Soil-moisture sensors can help determine when irrigation is required, while weather stations can provide information about rainfall and evapotranspiration. Remote sensing and drones can identify crop stress and differences in crop growth across fields.

Precision nutrient management can use soil maps, crop sensors and yield maps to determine nutrient requirements for different parts of a field. Variable-rate technology can then apply fertilizers according to site-specific requirements.

The integration of sensors, remote sensing, artificial intelligence and mobile-based decision-support systems can further improve irrigation and fertilizer scheduling. These technologies can help farmers move from fixed management schedules towards **demand-based resource management.**

Benefits for Farmers and the Environment

Integrated nutrient and water management can provide both economic and environmental benefits. Efficient fertilizer application reduces unnecessary input expenditure, while efficient irrigation reduces pumping costs and water consumption. Improved nutrient-use efficiency can also reduce losses to water bodies and the atmosphere.

Maintaining soil organic matter improves soil health and may increase the long-term productivity of agricultural land. Efficient use of water can help reduce pressure on groundwater resources. Thus, INWM can contribute to sustainable intensification by improving crop productivity without proportionally increasing the consumption of natural resources.

Challenges and Future Prospects

The adoption of integrated nutrient and water management may be limited by lack of soil-testing facilities, inadequate irrigation infrastructure, limited farmer awareness, high initial costs of efficient irrigation systems and insufficient technical guidance. Small and fragmented landholdings can also make the adoption of certain technologies difficult.

Extension services and farmer training are therefore essential. Demonstration plots can help farmers understand the economic and agronomic benefits of improved nutrient and water management. Custom-hiring and service-provider models can make technologies such as drip irrigation, soil sensors and precision fertilizer application more accessible.

In the future, integration of digital agriculture, artificial intelligence, remote sensing and automated irrigation systems can make nutrient and water management increasingly precise. However, technology

should complement rather than replace basic agronomic principles such as soil testing, balanced fertilization, proper irrigation scheduling and soil-moisture conservation.

Conclusion

Integrated Nutrient and Water Management is an important approach for achieving productive, profitable and sustainable agriculture. By combining balanced nutrient supply, organic inputs, efficient irrigation, moisture conservation and scientific scheduling, farmers can improve crop productivity while reducing wastage of valuable resources. Integration of modern technologies with sound agronomic practices can further improve nutrient-use efficiency and water-use efficiency. Wider adoption of INWM can contribute to improved soil health, conservation of water resources, reduced production costs and greater resilience of agricultural systems.

References

1. Dobermann, A. and Cassman, K.G. 2002. Plant nutrient management for enhanced productivity in intensive grain production systems of the United States and Asia. *Plant and Soil*, **247**: 153–175.
2. FAO. 2017. *Climate-Smart Agriculture Sourcebook*. 2nd ed. Food and Agriculture Organization of the United Nations, Rome.

3. Fixen, P.E., Brentrup, F., Bruulsema, T.W., Garcia, F., Norton, R. and Zingore, S. 2015. Nutrient/fertilizer use efficiency: Measurement, current situation and trends. In: *Managing Water and Fertilizer for Sustainable Agricultural Intensification*. IFA, Paris.
4. Havlin, J.L., Tisdale, S.L., Nelson, W.L. and Beaton, J.D. 2014. *Soil Fertility and Fertilizers: An Introduction to Nutrient Management*. 8th ed. Pearson, New Delhi.
5. Howell, T.A. 2001. Enhancing water use efficiency in irrigated agriculture. *Agronomy Journal*, **93**: 281–289.
6. Rockström, J., Karlberg, L., Wani, S.P., Barron, J., Hatibu, N., Oweis, T., Bruggeman, A., Farahani, J. and Qiang, Z. 2010. Managing water in rainfed agriculture—The need for a paradigm shift. *Agricultural Water Management*, **97**: 543–550.
7. Stewart, B.A. and Lal, R. 2018. *Advances in Soil and Water Conservation*. CRC Press, Boca Raton.
8. Vanlauwe, B., Wendt, J., Giller, K.E., Corbeels, M., Gerard, B. and Nolte, C. 2014. A fourth principle is required to define Conservation Agriculture in sub-Saharan Africa: The appropriate use of fertilizer to enhance crop productivity. *Field Crops Research*, **155**: 10–13.