NEW ERA AGRICULTURE MAGAZINE

Chromosomes: The Hidden Architects of Crop Improvement

Apeksha M. Suryawanshi[1], Kiran N. More[2], Shrikrishna B. Narale[3]

Introduction:

Behind every high-yielding, disease-resistant and climate-resilient crop lies an invisible blueprint—the **chromosome**. Although chromosomes cannot be seen with the naked eye, they carry the genetic information that controls virtually every characteristic of a plant, from plant height and flowering time to grain quality, disease resistance and stress tolerance. For plant breeders, chromosomes are therefore much more than structures visible under a microscope; they are the **architectural framework of crop improvement**.

Modern agriculture faces enormous challenges. Rapidly changing climates, emerging diseases, declining genetic diversity and increasing demand for food require crops that can perform reliably under difficult conditions. To develop such crops, breeders need to understand not only individual genes but also how genes are organized, inherited and exchanged through chromosomes.

What Exactly Are Chromosomes?

Chromosomes are thread-like structures present in the nucleus of plant cells. They are composed mainly of **DNA associated with proteins**, particularly histones. DNA contains genes—the functional units of heredity—that provide instructions for producing proteins and regulating biological processes. The number of chromosomes differs among crop species. For example, **bread wheat possesses 42 chromosomes (2n = 42)**, while maize has **20 chromosomes (2n = 20)** and rice has **24 chromosomes (2n = 24)**. Each chromosome contains thousands of genes arranged in a specific order.

Interestingly, chromosome number alone does not determine the complexity or usefulness of a crop. What matters greatly is **which genes are present, how they are organized and how they interact**.

Chromosomes and Crop Diversity

Genetic variation is the raw material of plant breeding. During meiosis, chromosomes undergo **recombination**, allowing genetic

Apeksha M. Suryawanshi[1], Kiran N. More[2], Shrikrishna B. Narale[3]
[1] Department of Animal Husbandry and Dairy Science (AHDS), Mahatma Phule Krishi Vidyapeeth (MPKV), Rahuri, Maharashtra, India.
[2] Ph.D. Scholar, Department of Agronomy, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth (DBSKKV), Dapoli, Maharashtra, India.
[3] Ph.D. Scholar, Department of Biochemistry, Mahatma Phule Krishi Vidyapeeth (MPKV), Rahuri, Maharashtra, India.

information from parents to be reshuffled in their offspring. This natural process creates new combinations of genes that breeders can select for desirable characteristics. For example, a breeder may cross a high-yielding variety with a disease-resistant donor. Through successive generations of selection, desirable chromosome segments from both parents can be combined into a superior breeding line.

Thus, every new variety represents a carefully selected combination of chromosome segments.

The Power of Polyploidy

One of the most fascinating features of plant chromosomes is **polyploidy**, in which plants possess more than two complete sets of chromosomes.Many important crops have evolved through polyploidy. **Bread wheat is a classic example**, possessing three related genomes—A, B and D—with seven chromosomes in each genome.

Polyploidy can provide plants with additional genetic material, sometimes contributing to increased adaptability, larger organs and enhanced genetic flexibility. It has therefore played an enormous role in the evolution and diversification of cultivated plants.

Wild Relatives: Hidden Treasure Houses

Wild relatives of crops often contain valuable genes that have been lost or reduced during domestication. These genes may provide resistance to diseases, insects, drought, salinity, heat or other stresses.

But how can breeders identify and transfer these useful chromosome segments?

This is where **cytogenetics** becomes extremely powerful.

Techniques such as **fluorescence in situ hybridization (FISH)** and **genomic in situ hybridization (GISH)** allow researchers to visualize chromosomes and identify

Table 1. Chromosome Numbers of Important Crop Plants

Sr. No.	Crop	Scientific name	Chromosome number (2n)	Ploidy level
1	Bread wheat	*Triticum aestivum*	42	Hexaploid
2	Durum wheat	*Triticum durum*	28	Tetraploid
3	Rice	*Oryza sativa*	24	Diploid
4	Maize	*Zea mays*	20	Diploid
5	Barley	*Hordeum vulgare*	14	Diploid
6	Chickpea	*Cicer arietinum*	16	Diploid
7	Soybean	*Glycine max*	40	Diploid
8	Cotton (upland cotton)	*Gossypium hirsutum*	52	Allotetraploid
9	Potato	*Solanum tuberosum*	48	Tetraploid
10	Groundnut	*Arachis hypogaea*	40	Allotetraploid

chromosome segments originating from different species. Imagine chromosomes as coloured threads. FISH and GISH can help scientists determine **which thread came from which parent or species**. This makes it possible to track valuable alien chromosome segments during breeding.

Chromosome Engineering: Designing Better Crops

Traditional crossing is not always sufficient to transfer a desirable gene from a wild relative. Sometimes unwanted genes are transferred along with the desired one. This phenomenon is known as **linkage drag**.

Chromosome engineering provides strategies to overcome this challenge.

Through controlled crossing, recombination and cytogenetic screening, breeders can attempt to reduce the size of unwanted chromosome segments while retaining the useful gene. This approach has been particularly important in **wheat improvement**, where chromosome segments from related species have been introduced to improve resistance to diseases and other stresses.

From Microscope to Molecular Markers

Cytogenetics has evolved dramatically.

Earlier, researchers mainly studied chromosome number, size, shape and pairing under a microscope. Today, cytogenetics works together with **molecular markers, DNA sequencing, genomics and bioinformatics**. A chromosome region identified through cytogenetic analysis can now be investigated using molecular markers and genome sequences. Researchers can locate genes and quantitative trait loci (QTLs), develop markers closely linked to useful traits and accelerate their transfer into elite varieties. This integration has created a powerful bridge:

Chromosome → DNA → Gene → Marker → Trait → Improved Variety

Chromosomes Meet Genome Editing

The arrival of **CRISPR-based genome editing** has taken crop improvement to another level. Genome editing can make precise changes in selected genes, while cytogenetics provides the broader chromosome-level context.

However, genome editing does not replace cytogenetics. Chromosome number, ploidy, chromosome structure and genome stability remain important considerations, particularly in polyploid crops such as wheat. The future of breeding is therefore likely to involve an integration of **cytogenetics + genomics + molecular breeding + genome editing**.

A New Era of Chromosome-Based Breeding

Modern plant breeding is moving from simply selecting plants with desirable appearances to understanding the genetic architecture underlying those traits.

Advanced technologies can now help breeders visualize chromosomes, sequence genomes, identify QTLs, detect structural variations and track useful chromosome segments with remarkable precision. This knowledge is especially important for developing crops capable of withstanding **drought, heat, salinity, emerging diseases and unpredictable climatic conditions**.

Conclusion

Chromosomes may be microscopic, but their influence on agriculture is enormous. They carry the genetic instructions that shape crops and provide the framework through which genetic variation is inherited and reshuffled. From the discovery of chromosome numbers to modern FISH, GISH, molecular markers, genome sequencing and CRISPR, our

Table 2. Major Cytogenetic Tools and Their Applications in Crop Improvement

Cytogenetic tool	What does it do?	Application in plant breeding
Karyotyping	Examines chromosome number, size and morphology	Identifies chromosomal abnormalities and ploidy differences
FISH (Fluorescence In Situ Hybridization)	Uses fluorescent probes to locate specific DNA sequences on chromosomes	Identifies chromosome regions and helps detect structural changes
GISH (Genomic In Situ Hybridization)	Distinguishes chromosomes or chromosome segments from different species	Tracks alien chromosome introgressions from wild relatives
Chromosome banding	Reveals characteristic chromosome patterns	Helps identify individual chromosomes and structural rearrangements
Meiotic cytology	Examines chromosome pairing and segregation during meiosis	Assesses chromosome behaviour, fertility and genome stability
Chromosome engineering	Manipulates chromosome segments through breeding and other techniques	Transfers useful genes from wild relatives into cultivated crops
Molecular cytogenetics	Combines cytogenetic methods with molecular tools	Supports chromosome identification and the localization of useful genetic regions
Chromosome-scale genome analysis	Integrates chromosome information with DNA sequencing and genome assembly	Facilitates gene discovery, structural variation analysis and precision breeding

understanding of chromosomes has transformed plant breeding. The next generation of crops will not be improved by looking only at the plant above the soil. Breeders will increasingly look **inside the cell—to the chromosomes and the genomes they carry**.

The future of crop improvement may be written in DNA, but its architecture is built on chromosomes.

Refrences

1. Boehm, J., Jr., & Cai, X. (2024). Enrichment and diversification of the wheat genome via alien introgression. *Plants, 13*(3), Article 339. Full-text article
2. Chen, Q. (2005). Detection of alien chromatin introgression from *Thinopyrum* into wheat using S genomic DNA as a probe—a landmark approach for *Thinopyrum* genome research. *Cytogenetic and Genome Research, 109*(1–3), 350–359.
3. Friebe, B., Jiang, J., Raupp, W. J., McIntosh, R. A., & Gill, B. S. (1996). Characterization of wheat-alien translocations conferring resistance to diseases and pests: Current status. *Euphytica, 91*, 59–87.
4. Pauk, J. (2016). Alien introgression in wheat: Cytogenetics, molecular biology, and genomics. *Cereal Research Communications, 44*, 535. Publisher's article page
5. Qi, L. L., Friebe, B., Zhang, P., & Gill, B. S. (2007). Homoeologous recombination, chromosome engineering and crop improvement. *Chromosome Research, 15*, 3–19.
6. Qi, L. L., Pumphrey, M. O., Friebe, B., Chen, P. D., & Gill, B. S. (2008). Molecular cytogenetic characterization of alien introgressions with gene *Fhb3* for resistance to Fusarium head blight disease of wheat. *Theoretical and Applied Genetics, 117*, 1155–1166.
7. *Review of the application of modern cytogenetic methods (FISH/GISH) to the study of reticulation (polyploidy/hybridisation).* (2010). *Genes, 1*(2), 166–192. Full-text article
8. Younis, A., Ramzan, F., Hwang, Y. J., & Lim, K. B. (2015). FISH and GISH: Molecular cytogenetic tools and their applications in ornamental plants. *Plant Cell Reports, 34*(9), 1477–1488. Publisher's article page
9. Zhang, W., Cao, Y., Zhang, M., Zhu, X., Ren, S., Long, Y., Gyawali, Y., Chao, S., Xu, S. S., & Cai, X. (2017). Meiotic homoeologous recombination-based alien gene introgression in the genomics era of wheat. *Crop Science, 57*, 1189–1198.

10. Zhang, W., Zhang, M., Zhu, X., Cao, Y., Sun, Q., Ma, G., Chao, S., Yan, C., Xu, S. S., & Cai, X. (2018). Molecular cytogenetic and genomic analyses reveal new insights into the origin of the wheat B genome. *Theoretical and Applied Genetics, 131*, 365–375.

NEW ERA
AGRICULTURE MAGAZINE